

# First results from DAMA/LIBRA (and the combined analysis with DAMA/NaI)



P. Belli  
INFN-Roma Tor Vergata


XXth RENCONTRES DE BLOIS  
Blois, May 2008

# Roma2, Roma1, LNGS, IHEP/Beijing

- + by-products and small scale expts.: INR-Kiev
- + neutron meas.: ENEA-Frascati
- + in some studies on  $\beta\beta$  decays (DST-MAE project): IIT Kharagpur, India




## DAMA: an observatory for rare processes @LNGS





# Relic DM particles from primordial Universe



(& invisible axions,  $\nu$ 's)

&


## Right halo model and parameters?



# Some direct detection processes:

- Scatterings on nuclei

→ detection of nuclear recoil energy



- Inelastic Dark Matter:  $W + N \rightarrow W^* + N$

→ W has Two mass states  $\chi^+$ ,  $\chi^-$  with  $\delta$  mass splitting

→ Kinematical constraint for the inelastic scattering of  $\chi^-$  on a nucleus


$$\frac{1}{2} \mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

- Excitation of bound electrons in scatterings on nuclei

→ detection of recoil nuclei + e.m. radiation


- Conversion of particle into e.m. radiation

→ detection of  $\gamma$ , X-rays,  $e^-$



- Interaction only on atomic electrons


→ detection of e.m. radiation



- Interaction of light DMp (LDM) on  $e^-$  or nucleus with production of a lighter particle

→ detection of electron/nucleus recoil energy

e.g. sterile  $\nu$



e.g. signals from these candidates are **completely lost** in experiments based on "rejection procedures" of the e.m. component of their rate




... also see the Khlopov's talk yesterday

- ... and more

The direct detection experiments can be classified in two classes, depending on what they are based:

1. on the recognition of the signals due to Dark Matter particles with respect to the background by using a “model-independent” signature

2. on the use of uncertain techniques of rejection of electromagnetic background (adding systematical effects and lost of candidates with pure electromagnetic productions)






# The annual modulation: a model independent signature for the investigation of Dark Matter particles component in the galactic halo

With the present technology, the annual modulation is the main model independent signature for the DM signal. Although the modulation effect is expected to be relatively small **a suitable large-mass, low-radioactive set-up with an efficient control of the running conditions would point out its presence.**

Drukier, Freese, Spergel PRD86  
Freese et al. PRD88



- $v_{\text{sun}} \sim 232 \text{ km/s}$  (Sun velocity in the halo)
- $v_{\text{orb}} = 30 \text{ km/s}$  (Earth velocity around the Sun)
- $\gamma = \pi/3$
- $\omega = 2\pi/T$        $T = 1 \text{ year}$
- $t_0 = 2^{\text{nd}} \text{ June}$  (when  $v_{\oplus}$  is maximum)

$$v_{\oplus}(t) = v_{\text{sun}} + v_{\text{orb}} \cos\gamma \cos[\omega(t-t_0)]$$

$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k} \cos[\omega(t-t_0)]$$

**Expected rate in given energy bin changes because the annual motion of the Earth around the Sun moving in the Galaxy**


## Requirements of the annual modulation

- 1) Modulated rate according cosine
- 2) In a definite low energy range
- 3) With a proper period (1 year)
- 4) With proper phase (about 2 June)
- 5) For single hit events in a multi-detector set-up
- 6) With modulation amplitude in the region of maximal sensitivity must be  $< 7\%$  for usually adopted halo distributions, but it can be larger in case of some possible scenarios

To mimic this signature, spurious effects and side reactions must not only - obviously - be able to account for the whole observed modulation amplitude, but also to satisfy contemporaneously all the requirements

# The $\approx 100$ kg NaI(Tl) set-up (DAMA/NaI)

- 9 highly radiopure NaI(Tl) of 9,7 kg coupled through 10 cm long Tetrasil-B light guides to 2 low background PMTs (specially developed).
- Detectors enclosed in a sealed copper box, continuously maintained in HP Nitrogen atmosphere, in slightly overpressure with respect to the external environment.
- A suitable low background hard shield against e.m. and neutron background was realized using very high radiopure Cu and Pb bricks, Cd foils and 10/40 cm polyethylene/paraffin; the hard shield is also sealed in a plexiglas box and maintained in the high purity HP Nitrogen atmosphere.
- Etc.



Model independent evidence for the presence of DM particles in the galactic halo

**Performances:** N.Cim.A112(1999)545-575, EPJC18(2000)283, Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127

## Results on rare processes:

- Possible Pauli exclusion principle violation PLB408(1997)439
- CNC processes PRC60(1999)065501
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell) PLB460(1999)235
- Search for solar axions PLB515(2001)6
- Exotic Matter search EPJdirect C14(2002)1
- Search for superdense nuclear matter EPJA23(2005)7
- Search for heavy clusters decays EPJA24(2005)51


## Results on DM particles:

- PSD PLB389(1996)757
- Investigation on diurnal effect N.Cim.A112(1999)1541
- Exotic Dark Matter search PRL83(1999)4918
- Annual Modulation Signature  
PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512,  
PLB480(2000)23, EPJC18(2000)283, PLB509(2001)197,  
EPJC23(2002)61, PRD66(2002)043503, Riv.N.Cim.26 n.1 (2003)1,  
IJMPD13(2004)2127, IJMPA21(2006)1445, EPJC47(2006)263,  
IJMPA22(2007)3155, EPJC53(2008)205, PRD77(2008)023506 +  
arXiv:0802.4336, other works in progress ...

*data taking completed on July 2002, last data release 2003; still producing results on corollary quests for the candidate particle and the possible astrophysical, nuclear and particle physics scenarios*

**total exposure (7 annual cycles) 0.29 ton x yr**

# The model-independent evidence by DAMA/NaI well compatible with several candidates in several of the many astrophysical, nuclear and particle physics scenarios; other ones are open



Neutralino as LSP in SUSY theories

Various kinds of WIMP candidates with several different kind of interactions  
Pure SI, pure SD, mixed + Migdal effect + channeling, ... (from low to high mass)

WIMP with preferred inelastic scattering

Mirror Dark Matter

Light Dark Matter

a heavy  $\nu$  of the 4-th family

Pseudoscalar, scalar or mixed light bosons with axion-like interactions

Dark Matter (including some scenarios for WIMP) electron-interacting

Sterile neutrino


Self interacting Dark Matter

Elementary Black holes such as the Daemons

heavy exotic candidates, as "4th family atoms", ...

Kaluza Klein particles

... and more



Possible model dependent positive hints from indirect searches not in conflict with DAMA/NaI result  
(but interpretation, evidence itself, derived mass and cross sections depend e.g. on bckg modeling, on DM spatial velocity distribution in the galactic halo, etc.)

Available results from direct searches using different target materials and approaches and from searches for up-going muons do not give any robust conflict



# The new DAMA/LIBRA set-up ~250 kg NaI(Tl) (Large sodium Iodide Bulk for RARE processes)

As a result of a second generation R&D for more radiopure NaI(Tl)  
by exploiting new chemical/physical radiopurification techniques  
(all operations involving crystals and PMTs - including photos - in HP Nitrogen atmosphere)



installing DAMA/LIBRA detectors




assembling a DAMA/ LIBRA detector

detectors during installation; in the  
central and right up detectors the new  
shaped Cu shield surrounding light  
guides (acting also as optical windows)  
and PMTs was not yet applied



filling the inner Cu box with  
further shield



closing the Cu box  
housing the detectors



view at end of detectors'  
installation in the Cu box




# The DAMA/LIBRA set-up

For details, radiopurity, performances, procedures, etc.  
see [arXiv:0804:2738](https://arxiv.org/abs/0804.2738) to appear on NIMA

Polyethylene/  
paraffin

- 25 × 9.7 kg NaI(Tl) in a 5x5 matrix
- two Suprasil-B light guides directly coupled to each bare crystal
- two PMTs working in coincidence at the single ph. el. threshold

## Installation



- OFHC low radioactive copper
- Low radioactive lead
- Cadmium foils
- Polyethylene/Paraffin
- Concrete from GS rock




~ 1m concrete from GS rock

- Dismounting/Installing protocol (with "Scuba" system)
- All the materials selected for low radioactivity
- Multicomponent passive shield
- Three-level system to exclude Radon from the detectors
- Calibrations in the same running conditions as production runs
- Installation in air conditioning + huge heat capacity of shield
- Monitoring/alarm system; many parameters acquired with the production data
- Pulse shape recorded by Waweform Analyzer *TVS641A* (2chs per detector), 1 *Gsample/s*, 8 bit, bandwidth 250 MHz
- Data collected from low energy up to MeV region, despite the hardware optimization was done for the low energy



# Some on residual contaminants in new NaI(Tl) detectors



$\alpha/e$  pulse shape discrimination has practically 100% effectiveness in the MeV range

The measured  $\alpha$  yield in the new DAMA/LIBRA detectors ranges from 7 to some tens  $\alpha/\text{kg}/\text{keV}$

Second generation R&D for new DAMA/LIBRA crystals: new selected powders, physical/chemical radiopurification, new selection of overall materials, new protocol for growing and handling

## $^{232}\text{Th}$ residual contamination

From time-amplitude method. If  $^{232}\text{Th}$  chain at equilibrium: it ranges from 0.5 ppt to 7.5 ppt

## $^{238}\text{U}$ residual contamination

First estimate: considering the measured  $\alpha$  and  $^{232}\text{Th}$  activity, if  $^{238}\text{U}$  chain at equilibrium  $\Rightarrow$   $^{238}\text{U}$  contents in new detectors typically range from 0.7 to 10 ppt



$^{238}\text{U}$  chain splitted into 5 subchains:  $^{238}\text{U} \rightarrow ^{234}\text{U} \rightarrow ^{230}\text{Th} \rightarrow ^{226}\text{Ra} \rightarrow ^{210}\text{Pb} \rightarrow ^{206}\text{Pb}$

Thus, in this case:  $(2.1 \pm 0.1)$  ppt of  $^{232}\text{Th}$ ;  $(0.35 \pm 0.06)$  ppt for  $^{238}\text{U}$

and:  $(15.8 \pm 1.6)$   $\mu\text{Bq}/\text{kg}$  for  $^{234}\text{U} + ^{230}\text{Th}$ ;  $(21.7 \pm 1.1)$   $\mu\text{Bq}/\text{kg}$  for  $^{226}\text{Ra}$ ;  $(24.2 \pm 1.6)$   $\mu\text{Bq}/\text{kg}$  for  $^{210}\text{Pb}$ .

## $^{\text{nat}}\text{K}$ residual contamination

The analysis has given for the  $^{\text{nat}}\text{K}$  content in the crystals values not exceeding about 20 ppb




## $^{129}\text{I}$ and $^{210}\text{Pb}$

$^{129}\text{I}/^{\text{nat}}\text{I} \approx 1.7 \times 10^{-13}$  for all the new detectors

$^{210}\text{Pb}$  in the new detectors:  $(5 - 30)$   $\mu\text{Bq}/\text{kg}$ .

... more on arXiv:0804.2738


No sizeable surface pollution by Radon daughters, thanks to the new handling protocols









# DAMA/LIBRA: calibrations at low energy

Studied by using various external gamma sources ( $^{241}\text{Am}$ ,  $^{133}\text{Ba}$ ) and internal X-rays or gamma's ( $^{40}\text{K}$ ,  $^{125}\text{I}$ ,  $^{129}\text{I}$ )



The curves superimposed to the experimental data have been obtained by simulations

- **Internal  $^{40}\text{K}$ :** 3.2 keV due to X-rays/Auger electrons (tagged by 1461 keV  $\gamma$  in an adjacent detector).
- **Internal  $^{125}\text{I}$ :** 67.3 keV peak (EC from K shell + 35.5 keV  $\gamma$ ) and composite peak at 40.4 keV (EC from L,M,... shells + 35.5 keV  $\gamma$ ).
- **External  $^{241}\text{Am}$  source:** 59.5 keV  $\gamma$  peak and 30.4 keV composite peak.
- **External  $^{133}\text{Ba}$  source:** 81.0 keV  $\gamma$  peak.
- **Internal  $^{129}\text{I}$ :** 39.6 keV structure (39.6 keV  $\gamma$  +  $\beta$  spectrum).






$$\frac{\sigma_{LE}}{E} = \frac{(0.448 \pm 0.035)}{\sqrt{E(\text{keV})}} + (9.1 \pm 5.1) \cdot 10^{-3}$$

Routine calibrations with  $^{241}\text{Am}$

# DAMA/LIBRA: calibrations at high energy

The data are taken on the full energy scale up to the MeV region by means QADC's

Studied by using external sources of gamma rays (e.g.  $^{137}\text{Cs}$ ,  $^{60}\text{Co}$  and  $^{133}\text{Ba}$ ) and gamma rays of 1461 keV due to  $^{40}\text{K}$  decays in an adjacent detector, tagged by the 3.2 keV X-rays





$$\frac{\sigma_{HE}}{E} = \frac{(1.12 \pm 0.06)}{\sqrt{E(\text{keV})}} + (17 \pm 23) \cdot 10^{-4}$$


The signals (unlike low energy events) for high energy events are taken only from one PMT

# Noise rejection near the energy threshold

Typical pulse profiles of PMT noise and of scintillation event with the same area, just above the energy threshold of 2 keV



The different time characteristics of PMT noise (decay time of order of tens of ns) and of scintillation event (decay time about 240 ns) can be investigated building several variables



From the Waveform Analyser  
2048 ns time window:

$$X_1 = \frac{\text{Area (from 100 ns to 600 ns)}}{\text{Area (from 0 ns to 600 ns)}};$$

$$X_2 = \frac{\text{Area (from 0 ns to 50 ns)}}{\text{Area (from 0 ns to 600 ns)}}$$

- The separation between noise and scintillation pulses is very good.
- Very clean samples of scintillation events selected by stringent acceptance windows.
- The related efficiencies evaluated by calibrations with  $^{241}\text{Am}$  sources of suitable activity in the same experimental conditions and energy range as the production data (efficiency measurements performed each ~10 days; typically  $10^4$ - $10^5$  events per keV collected)

This is the only procedure applied to the analysed data



# Infos about DAMA/LIBRA data taking

arXiv:0804.2741

DAMA/LIBRA test runs: from March 2003 to September 2003

DAMA/LIBRA normal operation: from September 2003 to August 2004

High energy runs for TDs: September 2004  
to allow internal  $\alpha$ 's identification  
(approximative exposure  $\approx 5000 \text{ kg} \times \text{d}$ )

DAMA/LIBRA normal operation: from October 2004

## Data released here:

- four annual cycles:  $0.53 \text{ ton} \times \text{yr}$
- calibrations: acquired  $\approx 44 \text{ M}$  events from sources
- acceptance window eff: acquired  $\approx 2 \text{ M}$  events/keV

Period		Exposure (kg $\times$ day)	$\alpha - \beta^2$
DAMA/LIBRA-1	Sept. 9, 2003 - July 21, 2004	51405	0.562
DAMA/LIBRA-2	July 21, 2004 - Oct. 28, 2005	52597	0.467
DAMA/LIBRA-3	Oct. 28, 2005 - July 18, 2006	39445	0.591
DAMA/LIBRA-4	July 19, 2006 - July 17, 2007	49377	0.541
Total		192824 $\simeq 0.53 \text{ ton} \times \text{yr}$	0.537

DAMA/NaI (7 years) + DAMA/LIBRA (4 years)

total exposure:  $300555 \text{ kg} \times \text{day} = 0.82 \text{ ton} \times \text{yr}$

## Two remarks:

- One PMT problems after 6 months. Detector out of trigger since Sep. 2003 (it will be put again in operation at the 2008 upgrading)
- Residual cosmogenic  $^{125}\text{I}$  presence in the first year in some detectors (this motivates the Sept. 2003 as starting time)


**DAMA/LIBRA is  
continuously running**

# Cumulative low-energy distribution of the *single-hit* scintillation events

Single-hit events = each detector has all the others as anticoincidence


(Obviously differences among detectors are present depending e.g. on each specific level and location of residual contaminants, on the detector's location in the 5x5 matrix, etc.)

Efficiencies already accounted for



## About the energy threshold:


- The DAMA/LIBRA detectors have been calibrated down to the keV region. This assures a clear knowledge of the "physical" energy threshold of the experiment.
- It obviously profits of the relatively high number of available photoelectrons/keV (from 5.5 to 7.5).
- The two PMTs of each detector in DAMA/LIBRA work in coincidence with hardware threshold at single photoelectron level.
- Effective near-threshold-noise full rejection.
- The software energy threshold used by the experiment is 2 keV.



# Experimental *single-hit* residuals rate vs time and energy

- Model-independent investigation of the annual modulation signature has been carried out by exploiting the time behaviour of the residual rates of the *single-hit* events in the lowest energy regions of the DAMA/LIBRA data.
- These residual rates are calculated from the measured rate of the *single-hit* events (obviously corrections for the overall efficiency and for the acquisition dead time are already applied) after subtracting the constant part:

$$\left\langle r_{ijk} - flat_{jk} \right\rangle_{jk}$$




- $r_{ijk}$  is the rate in the considered  $i$ -th time interval for the  $j$ -th detector in the  $k$ -th energy bin
- $flat_{jk}$  is the rate of the  $j$ -th detector in the  $k$ -th energy bin averaged over the cycles.
- The average is made on all the detectors ( $j$  index) and on all the energy bins ( $k$  index)
- The weighted mean of the residuals must obviously be zero over one cycle.



# Model Independent Annual Modulation Result

DAMA/NaI (7 years) + DAMA/LIBRA (4 years) Total exposure: 300555 kg×day = 0.82 ton×yr  
 experimental single-hit residuals rate vs time and energy




2-4 keV

$$A = (0.0215 \pm 0.0026) \text{ cpd/kg/keV}$$

$$\chi^2/\text{dof} = 51.9/66 \quad \mathbf{8.3 \sigma \text{ C.L.}}$$

Absence of modulation? No

$$\chi^2/\text{dof} = 117.7/67 \Rightarrow P(A=0) = 1.3 \times 10^{-4}$$




2-5 keV

$$A = (0.0176 \pm 0.0020) \text{ cpd/kg/keV}$$

$$\chi^2/\text{dof} = 39.6/66 \quad \mathbf{8.8 \sigma \text{ C.L.}}$$

Absence of modulation? No

$$\chi^2/\text{dof} = 116.1/67 \Rightarrow P(A=0) = 1.9 \times 10^{-4}$$



2-6 keV

$$A = (0.0129 \pm 0.0016) \text{ cpd/kg/keV}$$

$$\chi^2/\text{dof} = 54.3/66 \quad \mathbf{8.2 \sigma \text{ C.L.}}$$

Absence of modulation? No

$$\chi^2/\text{dof} = 116.4/67 \Rightarrow P(A=0) = 1.8 \times 10^{-4}$$

The data favor the presence of a modulated behavior with proper features at  $8.2\sigma$  C.L.

# Model-independent residual rate for single-hit events

DAMA/NaI (7 years) + DAMA/LIBRA (4 years)

total exposure: 300555 kg×day = 0.82 ton×yr

Results of the fits keeping the parameters free:

	A (cpd/kg/keV)	T= $2\pi/\omega$ (yr)	t <sub>0</sub> (day)	C.L.
<b>DAMA/NaI (7 years)</b>				
(2÷4) keV	0.0252 ± 0.0050	1.01 ± 0.02	125 ± 30	5.0σ
(2÷5) keV	0.0215 ± 0.0039	1.01 ± 0.02	140 ± 30	5.5σ
(2÷6) keV	0.0200 ± 0.0032	1.00 ± 0.01	140 ± 22	6.3σ
<b>DAMA/LIBRA (4 years)</b>				
(2÷4) keV	0.0213 ± 0.0032	0.997 ± 0.002	139 ± 10	6.7σ
(2÷5) keV	0.0165 ± 0.0024	0.998 ± 0.002	143 ± 9	6.9σ
(2÷6) keV	0.0107 ± 0.0019	0.998 ± 0.003	144 ± 11	5.6σ
<b>DAMA/NaI + DAMA/LIBRA</b>				
(2÷4) keV	0.0223 ± 0.0027	0.996 ± 0.002	138 ± 7	8.3σ
(2÷5) keV	0.0178 ± 0.0020	0.998 ± 0.002	145 ± 7	8.9σ
(2÷6) keV	0.0131 ± 0.0016	0.998 ± 0.003	144 ± 8	8.2σ




# Modulation amplitudes, $A$ , of single year measured in the 11 one-year experiments of DAMA (NaI + LIBRA)

- The difference in the (2 – 6) keV modulation amplitudes between DAMA/NaI and DAMA/LIBRA depends mainly on the rate in the (5 – 6) keV energy bin.
- The modulation amplitudes for the (2 – 6) keV energy interval, obtained when fixing exactly the period at 1 yr and the phase at 152.5 days, are:  
(0.019  $\pm$  0.003) cpd/kg/keV for DAMA/NaI  
(0.011  $\pm$  0.002) cpd/kg/keV for DAMA/LIBRA.
- Thus, their difference: (0.008  $\pm$  0.004) cpd/kg/keV is  $\approx 2\sigma$  which corresponds to a modest, but non negligible probability.

Moreover:

The  $\chi^2$  test ( $\chi^2 = 4.9, 3.3$  and  $8.0$  over 10 d.o.f. for the three energy intervals, respectively) and the *run test* (lower tail probabilities of 74%, 61% and 11% for the three energy intervals, respectively) **accept at 90% C.L. the hypothesis that the modulation amplitudes are normally fluctuating around their best fit values.**



**Compatibility among the annual cycles**




# Power spectrum of single-hit residuals

(according to Ap.J.263(1982)835; Ap.J.338(1989)277)


Treatment of the experimental errors and time binning included here

**2-6 keV vs 6-14 keV**


DAMA/NaI (7 years)  
total exposure: 0.29 ton×yr



DAMA/LIBRA (4 years)  
total exposure: 0.53 ton×yr



DAMA/NaI (7 years) +  
DAMA/LIBRA (4 years)  
total exposure: 0.82 ton×yr



Principal mode in the 2-6 keV region:

DAMA/NaI  
 $2.737 \cdot 10^{-3} \text{ d}^{-1} \approx 1 \text{ y}^{-1}$

DAMA/LIBRA  
 $2.705 \times 10^{-3} \text{ d}^{-1} \approx 1 \text{ yr}^{-1}$

DAMA/NaI+LIBRA  
 $2.737 \times 10^{-3} \text{ d}^{-1} \approx 1 \text{ yr}^{-1}$


+

Not present in the 6-14 keV region (only aliasing peaks)

Clear annual modulation is evident in (2-6) keV while it is absence just above 6 keV

# Can a hypothetical background modulation account for the observed effect?

## • No Modulation above 6 keV



Mod. Ampl. (6-10 keV):  $(0.0016 \pm 0.0031)$ ,  $-(0.0010 \pm 0.0034)$ ,  $-(0.0001 \pm 0.0031)$  and  $-(0.0006 \pm 0.0029)$  cpd/kg/keV for DAMA/LIBRA-1, DAMA/LIBRA-2, DAMA/LIBRA-3, DAMA/LIBRA-4;  
 → they can be considered statistically consistent with zero

+

In the same energy region where the effect is observed: no modulation of the multiple-hits events (see next slide)

## • No modulation in the whole spectrum:

studying integral rate at higher energy, R90

- $R_{90}$  percentage variations with respect to their mean values for single crystal in the DAMA/LIBRA-1,2,3,4 running periods


→ cumulative gaussian behaviour with  $\sigma \approx 1\%$ , fully accounted by statistical considerations

- Fitting the behaviour with time, adding a term modulated according period and phase expected for Dark Matter particles:

consistent with zero

Period	Mod. Ampl.
DAMA/LIBRA-1	$-(0.05 \pm 0.19)$ cpd/kg
DAMA/LIBRA-2	$-(0.12 \pm 0.19)$ cpd/kg
DAMA/LIBRA-3	$-(0.13 \pm 0.18)$ cpd/kg
DAMA/LIBRA-4	$(0.15 \pm 0.17)$ cpd/kg

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region →  $R_{90} \sim \text{tens cpd/kg}$  →  $\sim 100 \sigma$  far away



**No modulation in the background:**  
 these results account for all sources of bckg (+ see later)


# Multiple-hits events in the region of the signal - DAMA/LIBRA 1-4

- Each detector has its own TDs read-out  
→ pulse profiles of multiple-hits events (multiplicity > 1) acquired (exposure: 0.53 ton×yr).
- The same hardware and software procedures as the ones followed for single-hit events

signals by Dark Matter particles  
do not belong to multiple-hits  
events, that is:

multiple-hits events = Dark Matter particles events "switched off"

Evidence of annual modulation with proper features as required by the DM annual modulation signature is present in the *single-hit* residuals, while it is absent in the *multiple-hits* residual rate.



This result offers an additional strong support for the presence of Dark Matter particles in the galactic halo further excluding any side effect either from hardware or from software procedures or from background




## Energy distribution of the modulation amplitudes, $S_m$ , for the total exposure

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)]$$

DAMA/NaI (7 years) + DAMA/LIBRA (4 years)

total exposure: 300555 kg×day = 0.82 ton×yr

here  $T=2\pi/\omega=1$  yr and  $t_0=152.5$  day



A clear modulation is present in the (2-6) keV energy interval, while  $S_m$  values compatible with zero are present just above

In fact, the  $S_m$  values in the (6-20) keV energy interval have random fluctuations around zero with  $\chi^2$  equal to 24.4 for 28 degrees of freedom

# Statistical distributions of the modulation amplitudes ( $S_m$ )

a)  $S_m$  values for each detector, each annual cycle and each considered energy bin (here 0.25 keV)


b)  $\langle S_m \rangle$  = mean values over the detectors and the annual cycles for each energy bin;  $\sigma$  = errors associated to each  $S_m$

**DAMA/LIBRA (4 years)**


total exposure: 0.53 ton $\times$ yr

Each panel refers to each detector separately; 64 entries = 16 energy bins in 2-6 keV energy interval  $\times$  4 DAMA/LIBRA annual cycles

2-6 keV



Standard deviations of the variable  
 $(S_m - \langle S_m \rangle) / \sigma$   
 for the DAMA/LIBRA detectors



$r.m.s. \approx 1$

$0.80 < r.m.s. < 1.16$

Individual  $S_m$  values follow a normal distribution since  $(S_m - \langle S_m \rangle) / \sigma$  is distributed as a Gaussian with a unitary standard deviation (r.m.s.)



$S_m$  statistically well distributed in all the detectors and annual cycles

# Statistical analyses about modulation amplitudes ( $S_m$ )


$$x = (S_m - \langle S_m \rangle) / \sigma,$$

$$\chi^2 = \sum x^2$$

$\chi^2/\text{d.o.f.}$  values of  $S_m$  distributions for each DAMA/LIBRA detector in the (2–6) keV energy interval for the four annual cycles.

DAMA/LIBRA (4 years)

total exposure: 0.53 ton×yr



The  $\chi^2/\text{d.o.f.}$  values range from 0.7 to 1.28 (64 d.o.f. = 16 energy bins  $\times$  4 annual cycles)  
 $\Rightarrow$  at 95% C.L. the observed annual modulation effect is well distributed in all the detectors.

- The mean value of the twenty-four points is 1.072, slightly larger than 1. Although this can be still ascribed to statistical fluctuations, let us ascribe it to a possible systematics.
- In this case, one would have an additional error of  $\leq 5 \times 10^{-4}$  cpd/kg/keV, if quadratically combined, or  $\leq 7 \times 10^{-5}$  cpd/kg/keV, if linearly combined, to the modulation amplitude measured in the (2 – 6) keV energy interval.
- This possible additional error ( $\leq 4.7\%$  or  $\leq 0.7\%$ , respectively, of the DAMA/LIBRA modulation amplitude) can be considered as an upper limit of possible systematic effects





# Is there a sinusoidal contribution in the signal? Phase $\neq 152.5$ day?

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)] + Z_m \sin[\omega(t - t_0)] = S_0 + Y_m \cos[\omega(t - t^*)]$$

For Dark Matter signals:

- $|Z_m| \ll |S_m| \approx |Y_m|$
- $\omega = 2\pi/T$
- $t^* \approx t_0 = 152.5d$
- $T = 1 \text{ year}$

Slight differences from 2<sup>nd</sup> June are expected in case of contributions from non thermalized DM components (as e.g. the SagDEG stream)



E (keV)	$S_m$ (cpd/kg/keV)	$Z_m$ (cpd/kg/keV)	$Y_m$ (cpd/kg/keV)	$t^*$ (day)
2-6	$0.0122 \pm 0.0016$	$-0.0019 \pm 0.0017$	$0.0123 \pm 0.0016$	$144.0 \pm 7.5$
6-14	$0.0005 \pm 0.0010$	$0.0011 \pm 0.0012$	$0.0012 \pm 0.0011$	--

The analysis at energies above 6 keV, the analysis of the multiple-hit events and the statistical considerations about  $S_m$  already exclude any sizeable presence of systematical effects.

Additional investigations on the stability parameters

Modulation amplitudes obtained by fitting the time behaviours of main running parameters, acquired with the production data, when including a DM-like modulation

Running conditions stable  
at a level better than 1%



	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4
Temperature	$-(0.0001 \pm 0.0061) \text{ }^{\circ}\text{C}$	$(0.0026 \pm 0.0086) \text{ }^{\circ}\text{C}$	$(0.001 \pm 0.015) \text{ }^{\circ}\text{C}$	$(0.0004 \pm 0.0047) \text{ }^{\circ}\text{C}$
Flux N <sub>2</sub>	$(0.13 \pm 0.22) \text{ l/h}$	$(0.10 \pm 0.25) \text{ l/h}$	$-(0.07 \pm 0.18) \text{ l/h}$	$-(0.05 \pm 0.24) \text{ l/h}$
Pressure	$(0.015 \pm 0.030) \text{ mbar}$	$-(0.013 \pm 0.025) \text{ mbar}$	$(0.022 \pm 0.027) \text{ mbar}$	$(0.0018 \pm 0.0074) \text{ mbar}$
Radon	$-(0.029 \pm 0.029) \text{ Bq/m}^3$	$-(0.030 \pm 0.027) \text{ Bq/m}^3$	$(0.015 \pm 0.029) \text{ Bq/m}^3$	$-(0.052 \pm 0.039) \text{ Bq/m}^3$
Hardware rate above single photoelectron	$-(0.20 \pm 0.18) \times 10^{-2} \text{ Hz}$	$(0.09 \pm 0.17) \times 10^{-2} \text{ Hz}$	$-(0.03 \pm 0.20) \times 10^{-2} \text{ Hz}$	$(0.15 \pm 0.15) \times 10^{-2} \text{ Hz}$

All the measured amplitudes well compatible with zero  
+none can account for the observed effect


(to mimic such signature, spurious effects and side reactions must not only be able to account for the whole observed modulation amplitude, but also simultaneously satisfy all the 6 requirements)

# Example of Stability Parameters: DAMA/LIBRA-1

$R_{Hj}$  = hardware rate of j-th detector above single photoelectron



Running conditions stable at level < 1%  
Parameters distributions



All amplitudes well compatible with zero  
+ no effect can mimic the annual modulation




# Temperature


- Detectors in Cu housings directly in contact with multi-ton shield  
→ huge heat capacity ( $\approx 10^6$  cal/ $^{\circ}$ C)
- Experimental installation continuously air conditioned (2 independent systems for redundancy)
- Operating T of the detectors continuously controlled

Amplitudes for annual modulation in the operating T of the detectors **well compatible with zero**

	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4
T ( $^{\circ}$ C)	$-(0.0001 \pm 0.0061)$	$(0.0026 \pm 0.0086)$	$(0.001 \pm 0.015)$	$(0.0004 \pm 0.0047)$



Distribution of the relative variations of the operating T of the detectors



Distribution of the root mean square values of the operating T within periods with the same calibration factors (typically  $\approx 7$  days):

mean value  $\approx 0.04^{\circ}$ C

Considering the slope of the light output  $\approx -0.2\%/^{\circ}$ C:  
relative light output variation  $< 10^{-4}$ :


$< 10^{-4}$  cpd/kg/keV ( $< 0.5\%$   $S_m^{\text{observed}}$ )

**An effect from temperature can be excluded**

**+ Any possible modulation due to temperature would always fail some of the peculiarities of the signature**


# Summarizing on a hypothetical background modulation in DAMA/LIBRA 1-4

- No Modulation above 6 keV




- No modulation in the whole energy spectrum

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region  $\rightarrow R_{90} \sim \text{tens cpd/kg} \rightarrow \sim 100 \sigma$  far away



- No modulation in the 2-6 keV *multiple-hits* residual rate



*multiple-hits* residual rate (green points) vs  
single-hit residual rate (red points)

No background modulation (and cannot mimic the signature):

all this accounts for the all possible sources of bckg

Nevertheless, additional investigations performed ...

# Can a possible thermal neutron modulation account for the observed effect?

**NO**


- Thermal neutrons flux measured at LNGS :

$$\Phi_n = 1.08 \cdot 10^{-6} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (N.Cim.A101(1989)959)}$$

- Experimental upper limit on the thermal neutrons flux “surviving” the neutron shield in DAMA/LIBRA:
  - studying triple coincidences able to give evidence for the possible presence of  $^{24}\text{Na}$  from neutron activation:

$$\Phi_n < 1.2 \times 10^{-7} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (90\%C.L.)}$$

- Two consistent upper limits on thermal neutron flux have been obtained with DAMA/NaI considering the same capture reactions and using different approaches.



Evaluation of the expected effect:

► Capture rate =  $\Phi_n \sigma_n N_T < 0.022 \text{ captures/day/kg}$

**HYPOTHESIS:** assuming very cautiously a 10% thermal neutron modulation:

⇒  $S_m^{(\text{thermal n})} < 0.8 \times 10^{-6} \text{ cpd/kg/keV} (< 0.01\% S_m^{\text{observed}})$

In all the cases of neutron captures ( $^{24}\text{Na}$ ,  $^{128}\text{I}$ , ...) a possible thermal n modulation induces a variation in all the energy spectrum


Already excluded also by  $R_{90}$  analysis

## MC simulation of the process

When  $\Phi_n = 10^{-6} \text{ n cm}^{-2} \text{ s}^{-1}$ :

$7 \cdot 10^{-5} \text{ cpd/kg/keV}$

$1.4 \cdot 10^{-3} \text{ cpd/kg/keV}$



E (MeV)

# Can a possible fast neutron modulation account for the observed effect?

NO

In the estimate of the possible effect of the neutron background cautiously not included the 1m concrete moderator, which almost completely surrounds (mostly outside the barrack) the passive shield

Measured fast neutron flux @ LNGS:

$$\Phi_n = 0.9 \cdot 10^{-7} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (Astropart.Phys.4 (1995)23)}$$

By MC: differential counting rate above 2 keV  $\approx 10^{-3}$  cpd/kg/keV

HYPOTHESIS: assuming - very cautiously - a 10% neutron modulation:  $\Rightarrow S_m^{(\text{fast n})} < 10^{-4} \text{ cpd/kg/keV} (< 0.5\% S_m^{\text{observed}})$

- Experimental upper limit on the fast neutrons flux “*surviving*” the neutron shield in DAMA/LIBRA:

- through the study of the inelastic reaction  $^{23}\text{Na}(n,n')^{23}\text{Na}^*(2076 \text{ keV})$  which produces two  $\gamma$ 's in coincidence (1636 keV and 440 keV):

$$\Phi_n < 2.2 \times 10^{-7} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (90\%C.L.)}$$

- well compatible with the measured values at LNGS. This further excludes any presence of a fast neutron flux in DAMA/LIBRA significantly larger than the measured ones.

Moreover, a possible fast n modulation would induce:

- ▶ a variation in all the energy spectrum (steady environmental fast neutrons always accompanied by thermalized component)

already excluded also by  $R_{90}$

- ▶ a modulation amplitude for multiple-hit events different from zero

already excluded by the multiple-hit events


Thus, a possible 5% neutron modulation (ICARUS TM03-01) cannot quantitatively contribute to the DAMA/NaI observed signal, even if the neutron flux would be assumed 100 times larger than measured by various authors over more than 15 years @ LNGS



# Radon

- Three-level system to exclude Radon from the detectors:
- Walls and floor of the inner installation sealed in Supronyl ( $2 \times 10^{-11} \text{ cm}^2/\text{s}$  permeability).
- Whole shield in plexiglas box maintained in HP Nitrogen atmosphere in slight overpressure with respect to environment
- Detectors in the inner Cu box in HP Nitrogen atmosphere in slight overpressure with respect to environment continuously since several years

measured values at level of sensitivity of the used radonmeter



Time behaviours of the environmental radon in the installation (i.e. after the Supronyl), from which in addition the detectors are excluded by other two levels of sealing!

Amplitudes for annual modulation of Radon external to the shield:

$\langle \text{flux} \rangle \approx 320 \text{ l/h}$

Over pressure  $\approx 3.1 \text{ mbar}$

	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4
Radon ( $\text{Bq/m}^3$ )	$-(0.029 \pm 0.029)$	$-(0.030 \pm 0.027)$	$(0.015 \pm 0.029)$	$-(0.052 \pm 0.039)$

NO DM-like modulation amplitude in the time behaviour of external Radon (from which the detectors are excluded), of HP Nitrogen flux and of Cu box pressure


## Investigation in the HP Nitrogen atmosphere of the Cu-box

- Study of the double coincidences of  $\gamma$ 's (609 & 1120 keV) from  $^{214}\text{Bi}$  Radon daughter
- Rn concentration in Cu-box atmosphere  $< 5.8 \cdot 10^{-2} \text{ Bq/m}^3$  (90% C.L.)
- By MC:  $< 2.5 \cdot 10^{-5} \text{ cpd/kg/keV}$  @ low energy for *single-hit* events (enlarged matrix of detectors and better filling of Cu box with respect to DAMA/NaI)
- An hypothetical 10% modulation of possible Rn in Cu-box:

$< 2.5 \times 10^{-6} \text{ cpd/kg/keV}$  ( $< 0.01\% S_m^{\text{observed}}$ )

An effect from Radon can be excluded

+ any possible modulation due to Radon would always fail some of the peculiarities of the signature and would affect also other energy regions



# Can the $\mu$ modulation measured by MACRO account for the observed effect?

Case of fast neutrons produced by muons

$$\begin{aligned}\Phi_\mu @ \text{LNGS} &\approx 20 \mu \text{ m}^{-2} \text{ d}^{-1} & (\pm 2\% \text{ modulated}) \\ \text{Neutron Yield @ LNGS: } Y &= 1 \div 7 \cdot 10^{-4} \text{ n } / \mu / (\text{g/cm}^2) & (\text{hep-ex/0006014}) \\ R_n = (\text{fast n by } \mu) / (\text{time unit}) &= \Phi_\mu Y M_{\text{eff}}\end{aligned}$$

Annual modulation amplitude at low energy due to  $\mu$  modulation:

where:

$$S_m^{(\mu)} = R_n g \varepsilon f_{\Delta E} f_{\text{single}} \cdot 2\% / (M_{\text{setup}} \Delta E)$$

$g$  = geometrical factor  
 $\varepsilon$  = detection efficiency by elastic scattering  
 $f_{\Delta E}$  = energy window ( $E > 2\text{keV}$ ) efficiency  
 $f_{\text{single}}$  = single hit efficiency

Hyp.:  $M_{\text{eff}} = 15 \text{ tons}$   
 $g \approx \varepsilon \approx f_{\Delta E} \approx f_{\text{single}} \approx 0.5$  (cautiously)  
Knowing that:  
 $M_{\text{setup}} \approx 250 \text{ kg}$  and  $\Delta E = 4\text{keV}$


$$S_m^{(\mu)} < (0.4 \div 3) \times 10^{-5} \text{ cpd/kg/keV}$$

**NO**

Moreover, this modulation also induces a variation in other parts of the energy spectrum  
It cannot mimic the signature: already excluded also by  $R_{90}$



# Noise

Distribution of variations of total hardware rates of the crystals over the single ph.el. threshold (that is from noise to “infinity”) during DAMA/LIBRA-1,2,3,4 running periods

cumulative gaussian behaviour fully accounted by expected statistical spread arising from the sampling time used for the rate evaluation

$R_{Hj}$  = hardware rate of j-th detector above single photoelectron  
 $\langle R_{Hj} \rangle$  = mean of  $R_{Hj}$  in the corresponding annual cycle

Amplitudes for annual modulation well compatible with zero:



	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4
Hardware rate (Hz)	$-(0.20 \pm 0.18) \times 10^{-2}$	$(0.09 \pm 0.17) \times 10^{-2}$	$-(0.03 \pm 0.20) \times 10^{-2}$	$(0.15 \pm 0.15) \times 10^{-2}$

## Can a noise tail account for the observed modulation effect?

Despite the good noise identification near energy threshold and the used very stringent acceptance window for scintillation events (this is only procedure applied to the data), the role of an hypothetical noise tail in the scintillation events has even been quantitatively investigated.

The modulation amplitude of the "Hardware Rate" (period and phase as for DM particles) is compatible with zero:

$$(0.03 \pm 0.09) \times 10^{-2} \text{ Hz} \longrightarrow < 1.8 \times 10^{-3} \text{ Hz (90\% CL)}$$

Hardware Rate = noise + bckg [up to  $\approx$  MeV] + signal [up to  $\approx$  6 keV]

- noise/crystal  $\approx 0.10$  Hz
- relative modulation amplitude from noise  $< 1.8 \times 10^{-3} \text{ Hz} / 2.5 \text{ Hz} \approx 7.2 \times 10^{-4}$  (90%CL)

even in the worst hypothetical case of 10% residual tail of noise in the data

relative modulation amplitude from noise at low energy  $< 7.2 \times 10^{-5}$

$< 10^{-4} \text{ cpd/kg/keV}$

**NO**

# The calibration factors

**DAMA/LIBRA-1,2,3,4**

- Distribution of the percentage variations ( $\varepsilon_{tdcal}$ ) of each energy scale factor ( $tdcal_k$ ) with respect to the value measured in the previous calibration ( $tdcal_{k-1}$ ) for the DAMA/LIBRA-1 to -4 annual cycles.
- Distribution of the percentage variations ( $\varepsilon_{HE}$ ) of the high energy scale factor with respect to the mean values for the DAMA/LIBRA-1 to -4 annual cycles.

$$\varepsilon_{tdcal} = \frac{tdcal_k - tdcal_{k-1}}{tdcal_{k-1}}$$

→ the low energy calibration factor for each detector is known with an uncertainty  $\ll 1\%$  during the data taking periods: **additional energy spread  $\sigma_{cal}$**


$$\sigma = \sqrt{\sigma_{res}^2 + \sigma_{cal}^2} \approx \sigma_{res} \cdot \left[ 1 + \frac{1}{2} \left( \frac{\sigma_{cal}}{\sigma_{res}} \right)^2 \right]; \frac{1}{2} \left( \frac{\sigma_{cal} / E}{\sigma_{res} / E} \right)^2 \leq 7.5 \cdot 10^{-4} \frac{E}{20keV}$$

**Negligible effect considering routine calibrations and energy resolution at low energy**

**Confirmation from MC: maximum relative contribution  $< 1 - 2 \times 10^{-4}$  cpd/kg/keV**

**No modulation in the energy scale  
+ cannot mimic the signature**


gaussian behaviours





# The efficiencies

2-8 keV



**Distribution of variations of the efficiency values with respect to their mean values during DAMA/LIBRA running periods**

**Time behaviour:** modulation amplitudes obtained by fitting the time behaviours of the efficiencies including a WIMP-like cosine modulation for DAMA/LIBRA running periods


Energy	Amplitudes ( $\times 10^{-3}$ )			
	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4
2-4 keV	$(0.3 \pm 0.6)$	$(0.1 \pm 0.6)$	$-(0.4 \pm 1.1)$	$-(0.4 \pm 1.0)$
4-6 keV	$(0.0 \pm 0.6)$	$-(0.7 \pm 0.6)$	$-(0.3 \pm 1.0)$	$-(0.7 \pm 1.0)$
6-8 keV	$-(0.3 \pm 0.6)$	$-(1.0 \pm 0.7)$	$-(0.2 \pm 0.8)$	$-(1.0 \pm 0.8)$
8-10 keV	$-(0.5 \pm 0.5)$	$-(0.5 \pm 0.5)$	$-(0.2 \pm 0.6)$	$(0.7 \pm 0.6)$

Energy	Modulation amplitudes (DAMA/LIBRA)
2-4 keV	$(0.1 \pm 0.4) \times 10^{-3}$
4-6 keV	$-(0.4 \pm 0.4) \times 10^{-3}$


**Amplitudes well compatible with zero  
+ cannot mimic the signature**

# Summary of the results obtained in the additional investigations of possible systematics or side reactions (DAMA/LIBRA - arXiv:0804.2741)

<i>Source</i>	<i>Main comment</i>	<i>Cautious upper limit (90% C.L.)</i>
<b>RADON</b>	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	$<2.5 \times 10^{-6}$ cpd/kg/keV
<b>TEMPERATURE</b>	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield→ huge heat capacity + T continuously recorded	$<10^{-4}$ cpd/kg/keV
<b>NOISE</b>	Effective full noise rejection near threshold	$<10^{-4}$ cpd/kg/keV
<b>ENERGY SCALE</b>	Routine + intrinsic calibrations	$<1-2 \times 10^{-4}$ cpd/kg/keV
<b>EFFICIENCIES</b>	Regularly measured by dedicated calibrations	$<10^{-4}$ cpd/kg/keV
<b>BACKGROUND</b>	No modulation above 6 keV; no modulation in the (2-6) keV <i>multiple-hits</i> events; this limit includes all possible sources of background	$<10^{-4}$ cpd/kg/keV
<b>SIDE REACTIONS</b>	Muon flux variation measured by MACRO	$<3 \times 10^{-5}$ cpd/kg/keV



+ even if larger they cannot satisfy all the requirements of annual modulation signature




Thus, they can not mimic the observed annual modulation effect

# ... about the interpretation of the direct DM experimental results

## The positive and model independent result of DAMA/NaI + DAMA/LIBRA

- Presence of modulation for 11 annual cycles at  $\sim 8.2\sigma$  C.L. with the proper distinctive features of the signature; all the features satisfied by the data over 11 independent experiments of 1 year each one
- Absence of known sources of possible systematics and side processes able to quantitatively account for the observed effect and to contemporaneously satisfy the many peculiarities of the signature



**No other experiment whose result can be directly compared in model independent way is available so far**




To investigate the nature and coupling with ordinary matter of the possible DM candidate(s), effective energy and time correlation analysis of the events has to be performed within given model frameworks

## Corollary quests for candidates

- astrophysical models:  $\rho_{DM}$ , velocity distribution and its parameters
- nuclear and particle Physics models
- experimental parameters

e.g. for WIMP class particles: SI, SD, mixed SI&SD, preferred inelastic, scaling laws on cross sections, form factors and related parameters, spin factors, halo models, etc.

- + different scenarios
- + multi-component halo?



**THUS**  
**uncertainties on models**  
**and comparisons**

- **In progress complete model dependent analyses** by applying maximum likelihood analysis in time and energy accounting for at least some of the many existing uncertainties in the field (as done by DAMA/NaI in Riv.N.Cim.26 n.1 (2003)1, IJMPD13(2004)2127, IJMPA21(2006)1445, EPJC47(2006)263, IJMPA22(2007)3155, EPJC53(2008)205, PRD77(2008)023506, arXiv:0802.4336 ), and to enlarge the investigations to other scenarios
- Just to offer some naive feeling on the complexity of the argument:


**experimental  $S_m$  values vs expected behaviours**

for some DM candidates in few of  
the many possible astrophysical,  
nuclear and particle physics  
scenarios and parameters values






**Examples** for few of the many possible scenarios superimposed to the measured modulation amplitudes  $S_{m,k}$



WIMP DM candidate (as in [4])  
 considering elastic scattering on nuclei  
 SI dominant coupling  
 $v_0 = 170 \text{ km/s}$




About the same C.L. ...scaling from NaI


**channeling contribution as in EPJC53(2008)205 considered for curve *b***

Curve label	Halo model (see ref. [4, 34])	Local density (GeV/cm <sup>3</sup> )	Set as in [4]	DM particle mass	$\xi\sigma_{SI}$ (pb)
<i>a</i>	A5 (NFW)	0.2	A	15 GeV	$3.1 \times 10^{-4}$
<i>b</i>	A5 (NFW)	0.2	A	15 GeV	$1.3 \times 10^{-5}$
<i>c</i>	A5 (NFW)	0.2	B	60 GeV	$5.5 \times 10^{-6}$
<i>d</i>	B3 (Evans power law)	0.17	B	100 GeV	$6.5 \times 10^{-6}$
<i>e</i>	B3 (Evans power law)	0.17	A	120 GeV	$1.3 \times 10^{-5}$

# Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$



WIMP DM candidate (as in [4])  
 Elastic scattering on nuclei  
 SI & SD mixed coupling  
 $v_0 = 170$  km/s




About the same C.L. ...scaling from NaI



$\theta = 2.435$

Curve label	Halo model (see ref. [4, 34])	Local density (GeV/cm <sup>3</sup> )	Set as in [4]	DM particle mass	$\xi\sigma_{SI}$ (pb)	$\xi\sigma_{SD}$ (pb)
<i>f</i>	A5 (NFW)	0.2	A	15 GeV	$10^{-7}$	2.6
<i>g</i>	A5 (NFW)	0.2	A	15 GeV	$1.4 \times 10^{-4}$	1.4
<i>h</i>	A5 (NFW)	0.2	B	60 GeV	$10^{-7}$	1.4
<i>i</i>	A5 (NFW)	0.2	B	60 GeV	$8.7 \times 10^{-6}$	$8.7 \times 10^{-2}$
<i>j</i>	B3 (Evans power law)	0.17	A	100 GeV	$10^{-7}$	1.7
<i>k</i>	B3 (Evans power law)	0.17	A	100 GeV	$1.1 \times 10^{-5}$	0.11



# Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$



**LDM candidate**  
(as in arXiv:0802.4336):  
inelastic interaction  
with electron or nucleus targets



**Light bosonic candidate**  
(as in IJMPA21(2006)1445):  
axion-like particles totally  
absorbed by target material



About the same C.L.

(NFW) halo model as in [4, 34], local density = 0.17 GeV/cm<sup>3</sup>, local velocity = 170 km/s

Curve label	DM particle	Interaction	Set as in [4]	$m_H$	$\Delta$	Cross section (pb)
$l$	LDM	coherent on nuclei	A	30 MeV	18 MeV	$\xi\sigma_m^{coh} = 1.8 \times 10^{-6}$
$m$	LDM	coherent on nuclei	A	100 MeV	55 MeV	$\xi\sigma_m^{coh} = 2.8 \times 10^{-6}$
$n$	LDM	incoherent on nuclei	A	30 MeV	3 MeV	$\xi\sigma_m^{inc} = 2.2 \times 10^{-2}$
$o$	LDM	incoherent on nuclei	A	100 MeV	55 MeV	$\xi\sigma_m^{inc} = 4.6 \times 10^{-2}$
$p$	LDM	coherent on nuclei	A	28 MeV	28 MeV	$\xi\sigma_m^{coh} = 1.6 \times 10^{-6}$
$q$	LDM	incoherent on nuclei	A	88 MeV	88 MeV	$\xi\sigma_m^{inc} = 4.1 \times 10^{-2}$
$r$	LDM	on electrons	–	60 keV	60 keV	$\xi\sigma_m^e = 0.3 \times 10^{-6}$

curve  $r$ : also pseudoscalar axion-like candidates (e.g. majoron)  
 $m_a=3.2$  keV  $g_{aee}=3.9 \cdot 10^{-11}$

# Conclusions

- DAMA/LIBRA over 4 annual cycles ( $0.53 \text{ ton} \times \text{yr}$ ) confirms the results of DAMA/NaI ( $0.29 \text{ ton} \times \text{yr}$ )

- The cumulative confidence level for the model independent evidence for presence of DM particle in the galactic halo is  $8.2 \sigma$  (total exposure  $0.82 \text{ ton} \times \text{yr}$ )
- The updating of corollary analyses in some of the many possible scenarios for DM candidates, interactions, halo models, nuclear/atomic properties, etc. is in progress. Further ones are under consideration also on the basis of literature



- Upgrading of the experimental set-up prepared and soon being performed in 2008
- Analyses/data taking to investigate other rare processes in progress/foreseen

Starting new data taking cycles after upgrading to improve the investigation, to disentangle at least some of the many possibilities, to investigate other features of DM particle component(s) and second order effects, etc..

A possible highly radiopure NaI(Tl) multi-purpose set-up DAMA/1 ton (proposed by DAMA in 1996) is at present at R&D phase

to deep investigate Dark Matter phenomenology at galactic scale

Interesting complementary information from accelerators and indirect searches in space are also expected soon...

